

Original article

Work-related musculoskeletal disorders among dental professionals: An evidence-based update

Senthil P. Kumar*, Vijaya Kumar*, Mohan Baliga**

ABSTRACT

Background: Numerous studies have reported prevalence rates for work-related musculoskeletal disorders (WRMSD) among dental professionals (DP).

Objective: The objective of this paper is to update the dental professionals, physical therapists, occupational therapists, epidemiologists and ergonomics specialists with the growing evidence on WRMSD in DP through a systematic literature search.

Materials and methods: Systematic review of studies in MEDLINE from 1972-2010 was conducted using search terms dental OR "dentist" work-related," "musculoskeletal" in title abstract for English articles. Consensus was obtained after independent blinded search and overall findings were summarized.

Main findings: Thirty-four included articles were heterogenous and studied all categories of DP which found one-year prevalence for MSK pain was from 24-100%. Neck and shoulder pain were the highest prevalent MSK symptom. Most of the reported symptoms were chronic and recurrent.

Conclusion: There was an overall very high prevalence of MSK pain among DP. Among DP, dental hygienists and dental students were more affected with MSK symptoms. The risk factors highly predictive of developing MSK pain were biopsychosocial, And further research on prevalence in Indian dental professionals is warranted before implementation of preventive educational programmes among dentists.

Key words: Occupational disorders; prevalence; public health; dental education

INTRODUCTION

Oral healthcare providers (OHP) or dental professionals (DP) include dentists, dental assistants, dental hygienists, dental students, dental nurses, dental laboratory workers, dental technicians and dental office workers[1]. Work-related musculoskeletal

disorders (WRMSD) are problems of the musculoskeletal system that significantly cause workplace problems, thus affecting occupational health, productivity and career of the working population[2]. Like all other professionals, DP are exposed to occupational health hazards which predispose them to develop a multitude of health problems.

A study by Jacobsen et al[3]in their survey of dental laboratory technicians found biannual period prevalence of health problems to be 79%. Among those with health problems, musculoskeletal problems were present among 68%, dermal (34%), respiratory (31%), neurological (26%), systemic (19%) and eyesight/hearing problems (15%). The study demonstrated the magnitude of musculoskeletal problems in comparison to

Author's Affiliation: *PT., (PhD)., Associate professor, Dept of Physiotherapy, Kasturba Medical College (Manipal University), Mangalore, India. **MDS., Professor and Head, Dept of Oral and Maxillofacial Surgery, Manipal College of Dental Sciences (Manipal University), Mangalore, India.

Reprints Requests: Dr. Vijay Kumar, PT., (PhD)., Associate professor, Dept of Physiotherapy, Kasturba Medical College (Manipal University), Mangalore, India.

(Received on 31.07.2011, accepted on 26.08.2011)

other health problems among DP. Another study by Murtomaa[4] in a survey of 68 dentists and 90 dental assistants found report of occupational disorders among 31% of dentists and 12% of dental assistants. Among them, 31% of affected dentists and 10% of affected assistants reported having been on sick leave. Most common disorders were back and neck musculoskeletal disorders, stress and allergic rashes.

Though other authors reported risk factors and associated prevalence for non-musculoskeletal disorders[5-9] among DP, the magnitude and proportion of symptoms due to musculoskeletal disorders is so huge that it has an enormous impact on the quality of life of DP.

OBJECTIVES

The objective of this paper is to update the dental professionals, physical therapists, occupational therapists, epidemiologists and ergonomics specialists with the growing evidence on WRMSD in DP through a systematic literature search.

METHODS

Independent search was conducted by two testers independently, using search terms “dental” “dentist” “work-related” “musculoskeletal” in title in National Library of Medicine, MEDLINE ,(www.pubmed.com) website. The retrieved citations and articles were appraised and mutual consensus was obtained in presence of the third tester. The included articles were then grouped qualitatively under overall prevalence, region of involvement, risk factors, impact, and treatments taken for WRMSD among DP.

RESULTS

Our search yielded 55 citations from 1972 to 2010. Of the 55 articles, we excluded non-English articles (12 papers), letters to editor (4 papers) or commentary articles (5 papers). Thus final number of included articles was 34.

Main findings

1)Prevalence

There were twelve studies on prevalence of MSK symptoms among DP.

Dental professionals

Four studies evaluated the prevalence of musculoskeletal (MSK) pain in general among dental professionals as a single group that comprised of two or more categories of professionals in dental healthcare.

Dajpratham et al[10] performed their survey on 164 dental personnel and, surprisingly ,high number of 158 participants (96.3%) reported presence of musculoskeletal pain. Of those with pain, 32 (20.3%) were clinical instructors, 52 (32.9%) were post-graduate students and 74 (46.8%) were dental assistants. Hayes et al¹¹ in their review, found that overall prevalence of musculoskeletal pain among dental professionals (dentists, dental hygienists and dental students) was 64-93%.

Sartorio et al[12] in their review, found a prevalence of work-related musculoskeletal disorders ranging from 54-93% among dental professionals (dentists, dental hygienists, dental auxiliaries). Lalumandier et al[13] surveyed 5,000 dental personnel, dentists and dental auxiliaries and found prevalence of at least one MSK symptom in the previous year among all the three groups.

Dental hygienists

Two studies evaluated prevalence of MSK among dental hygienists population. Anton et al¹⁴ in their survey of 95 dental hygienists found that at least one MSD was reported by 93% of the studied sample. Prevalence of carpal tunnel syndrome symptoms was 42%. Osborn et al¹⁵ found among 493 dental hygienists that 68% had MSK pain in eight bodily locations during previous year.

Dental students

Five studies evaluated prevalence among dental student population. de Carvalho et al¹⁶ in their study on 227 Brazilian dental students found an overall prevalence of musculoskeletal pain during or after clinical work in 173 students to (76.2%). Thornton et al[17] in their survey of 590 dental students reported one-

year prevalence of musculoskeletal symptoms among 61% which they felt during their clinical work. Abou-Atme *et al*[18] found in their five-year follow-up of dental students that temporomandibular joint (TMJ) pain, TMJ sounds, upper back pain and lower back pain had generally increasing trends from the first and second years to the third year and then a trend towards decrease in the fourth and fifth years.

Tezel *et al*[19] found prevalence of neuropathic symptoms co-existing with MSK pain among left- and right- handed dental students: weakness (42%, 40%), pins and needles (35%, 22%) and numbness (23%, 19%). Melis *et al*[20] in their survey of 392 dental students found greatest prevalence of headache followed by lower back pain versus a control comparison group of 114 psychology students.

Influence of dental work category

One study compared the prevalence rates between different categories of dental professionals. Rice *et al*²¹ found that among three categories of 45 dental workers (overall prevalence of 75.6%)- dentists, dental assistants and dental hygienists - the dental hygienists had greater prevalence of MSK symptoms and were at greater risk for developing upper extremity symptoms, carpal tunnel syndrome (CTS) and back pain.

2)Region of musculoskeletal pain

Nine studies evaluated the prevalence of MSK symptoms according to region of involvement among DP. Dajpratham *et al*[10] found shoulder pain in 114 (72.2%), neck pain in 111 (70.3%) and low back pain in 80 (50.6%) of the total 158 participants.

Hayes *et al*[22] (2009a) found neck pain (64.29%), lower back pain (57.94%) and shoulder pain (48.41%) prevalent among Australian undergraduate dental hygiene students.

Hayes *et al*[11] in their review found most prevalent regions for pain in dentists to be back (36.3-60.1%) and neck (19.8-85%) and for dental hygienists, the wrist and hand pain was most prevalent at 60-69.5%.

Samotoi *et al*[23] found most commonly reported problems in previous year among 323 dental therapists. They found prevalence of neck (56.8%), lower back (54%) and shoulder pain (52.4%) among their study population. Thornton *et al*[17] found that among those students who had musculoskeletal symptoms, neck pain represented 48%, shoulder 31%, back 44% and hands 20%. Tezel *et al*¹⁹ found left- and right-handed students had shoulder pain (78%, 58%), neck pain (67%, 43%), back pain (56%, 43%) upper limb pain (46%, 43%) and headaches (34%, 22%) respectively.

Finsen *et al*[24] reported one-year prevalence of 65% for neck/shoulder pain and 59% for low back disorders among dentists. Augustson and Morken[25] found one-year overall prevalence of 81% for MSK symptoms among 329 dentists with shoulder pain (45%), neck pain (47%), low back pain (49%), hand/wrist pain (21%), upper back pain (20%), hip pain (18%), knee pain (14%), elbow pain (12%) and ankle pain (10%) complaints. Oberg and Oberg[26] (1993) found that 62% had neck complaints and 815 had one/both shoulder pain in the previous 12 months among 38 dental hygienists.

Neck and shoulder symptoms

Three studies specifically evaluated prevalence characteristics for neck/shoulder pain in DP. Morse *et al*[27] reviewed studies on prevalence for neck and shoulder disorders among dental practitioners (dentists, dental hygienists and dental assistants). Dentists reported 26-73% neck symptoms and 20-65% for shoulder symptoms on a one-year period prevalence. Dental hygienists reported higher rates (54-83%) for neck and 35-76% for shoulder, and dental assistants at 38-62% and 27-62% for neck and shoulder respectively. Morse *et al*[28] found that neck symptoms were reported by 37%, 43% and 72% and shoulder symptoms by 11%, 20% and 35% respectively by students, students/assistants and experienced dental hygienists. Akesson *et al*[29] found pain ratings and Nordic questionnaire were sensitive and specific for detecting work-related neck and shoulder complaints among 90 female dental personnel surveyed.

Risk factors

There were twelve studies finding association between physical, mental and social risk factors and reporting of MSK symptoms among DP. Eleven studies were on biological risk factors, five on psychological and eleven studies on social risk factors.

Biological- individual (physical)

Kihara[30] in a survey of dentists in private clinics found that prevalence of hand and arms, neck and shoulders and low back symptoms differed between three work postures commonly adopted by clinicians. Trunk lateral bending and rotation increased the low back extensor muscle EMG activity. Morse et al[27] reviewed and found static awkward postures involving isometric contractions of trapezius. Hayes et al[22] (2009a) found that lack of regular physical exercise was highly associated with lower back pain.

Valachi[31] reported higher risk of occupational injury and symptoms in women compared to their male counterparts. Yamalik[32] reported muscular imbalance, neuromuscular inhibition and previous pain are all risk factors towards development of future MSK symptoms in OHPs. Repeated unnatural, deviated or inadequate working postures and forceful hand movements were well identified risk factors.

Morse et al[28] found self-reported shoulder pain was strongly associated with working above shoulder height and neck symptoms with working positions involving a bent neck posture. Tezel et al[19] found no significant differences in MSK symptoms because of gender. The studied left-handed dental students, however, had significantly more MSK symptom prevalence than their right-handed counterparts. Oberg and Oberg[26] found that frequency of neck/shoulder symptoms was greater in the right handed dental hygienists.

Anton et al[14] found significant association of age and body mass index with carpal tunnel syndrome in dental hygienists. Ylipaa et al[33] found that dental hygienists under 42 years

of age had more MSK disorder prevalence. In Michalak-Turcotte study (2000), risk factors for WRMSD among dental hygienists are repetitive motion, pinch-grasp force, vibration and prolonged awkward postures.

Akesson et al[34] found high frequency of MSK disorders reflected the high demands of vision and fine precise hand movements and working with unsupported elevated arms during dental hygiene practice. The authors also found lack of association between MSK symptoms and blood levels of mercury and selenium, BMI and smoking habit. Moen and Bjorvatn[35] found that female dentists were more symptomatic than male dentist in MSK disorders among 139 surveyed dental personnel (dentists, dental auxiliaries and office workers).

Psychological- cognitive affective (mental):

Locker et al[36] in their survey of 1000 Canadian dental assistants on self-perceived levels of work-related stress found 38.8% reporting moderately stressful and 14.5% reporting extremely stressful. Forty-Three percent of those reporting stress also reported their intention to change jobs to another profession. Warren[37] found strong association between psychosocial stress and musculoskeletal disorder causation among dental practitioners and dental hygienist students.

Ylipaa et al[33] found work-family overload and work relations were associated with more MSK disorders among dental hygienists. Ylipaa et al³⁸ found active leisure and psychosocial work factors (mastery of work) were associated with increased odds of good health and well-being among 575 surveyed dental hygienists. Augustson and Morken[25] found positive association between perceived workload and shoulder discomfort.

Social- environmental or ergonomic

Hayes et al[22] found that 16-20hours of desk-based study by dental hygiene students was associated with neck pain and 6-10hours on computer lead to greater predisposition to shoulder pain and upper back pain. Samotoi et al[23] reported that less number of dental therapists working with dental assistants had

shoulder symptoms. Those working with assistants however, reported higher prevalence of hip/thigh symptoms.

Thornton et al[17] found that third and final year dental students showed significant relationship between occurrence of symptoms and equipment utilization, work efficiency and general health. Yamalik[32] reported inadequate equipment or workplace designs and inappropriate work patterns as particular risk factors for MSK symptoms.

Morse et al[28] reported higher supervisor support was associated with fewer dental hygienists reporting neck and/or shoulder symptoms. More et al[39] found that greater use of ultrasonic scalers and manual instruments to around 8.2 hours per week and 10.1 hours per week respectively was associated with symptoms of numbness in upper extremities compared to 2.8 hours per week and 4.8 hours week for students without numbness symptoms.

Anton et al[14] found that the number of patients treated per day contribute as a significant risk factor for greater prevalence of CTS symptoms among dental hygienists. Ylipaa et al[33] found that dental hygienists practising in Australia are more associated with MSK disorders than their Swedish counterparts. Thus the authors' findings reiterated that the complex interdependent interaction between socio-demographic, occupational and cultural aspects influence a dental hygienist's work, health and well-being.

Ylipaa et al[38] found high clinical practice fraction, high management support increased the odds for good general health while work-family overload and high work efficiency decreased the odds. Increased work experience and scaling work duration per week was associated with finger disorders and upper body MSK disorders respectively. Ylipaa et al⁴⁰ found strainful ergonomics, solitary work, patient treatment hours and working hours were significant predictors for development of WRMSD and neck/shoulder pain among 364 dental hygienists. Those who worked in public dental health service experienced more symptoms than private practice.

Augustson and Morken[25] found sport activity participation was negatively associated with low back pain. Liss et al⁴¹ found that days worked per week, time with trunk rotated postures and years of experience were significant predictors of reported shoulder pain.

Impact or consequence of musculoskeletal pain

Two studies evaluated the impact of MSK pain on health care system. Dajpratham et al¹⁰ found usage of pain relieving medication (34.8%), seeking medical evaluation (32.3%), reduction in working hours (27.2%), difficulty sleeping (22.8%) and work absence (22.8%) among 158 participants with musculoskeletal symptoms. Osborn et al¹⁵ found 34% of dental hygienists with MSK pain reported that pain had affected their clinical practice, forcing them to practise fewer days, decreasing their endurance, reducing speed and quality, and/or altering operating positions.

Treatments taken for relief of musculoskeletal pain

Two studies evaluated the of treatments taken by DP with MSK pain. Dajpratham et al¹⁰ found Thai traditional massage (51.9%), medication (28.5%), physical therapy (15.8%), acupuncture (7.6%) and alternative medicine (4.4%) was taken for pain relief by their subjects. Augustson and Morken[25] found incorporating ergonomic equipment into dental practice alleviated shoulder discomfort.

Evidence for interventions

Ergonomic interventions

A study by Michalak-Turcotte[42] outlined the components of ergonomic evaluation and intervention program and another by Sanders and Turcotte[43] reported two cases of dental hygienists managed effectively.

DISCUSSION

This review found an overall prevalence estimate ranging from 23-100% in DP reporting from previous year at least one MSK

pain in a body region which was mostly neck and/or shoulder or upper back pain.

The involvement of upper body/neck/shoulder among DP is understandable from a biomechanical viewpoint. The work postures, use of equipment, work station and work demands necessitate a complex interplay of biopsychosocial variables which might have lead to increased reporting of symptoms.

These findings are of a wider range compared to the only other systematic review performed earlier by Hayes *et al*[11]which found a narrower range of prevalence rate (64-93%). The reasons could be attributable to the selection criteria (including both reviews and original articles,using comprehensive search terms and not using medical subject headings- MeSH) employed in this review and inclusion of studies on all categories of DP instead of only dentists and dental hygienists. The aim of this review was to provide a detailed and comprehensive data on the prevalence of MSK pain. Though inclusion of all DP caused the inevitable heterogeneity, the estimates are far beyond expectations even when all groups are combined and this indicates a huge public health issue.

Work place characteristics- the DP practise in a wide variety of settings such as academic, hospital, private practice, community centers, research centers influence not only the predisposition to MSK pain but also maintain the symptoms leading to chronicity and work-related disability. The risk factors for MSK pain in DP found in this review include not only biological but psychological and social as well. This adds support to existing evidence for multidisciplinary biopsychosocial rehabilitation as a treatment of choice for neck and shoulder pain among working age adult population[44].

One of the few limitations of this review was the subjective nature of collecting information from DP in the reviewed studies, using verbal screening questions rather than clinical examination. Some of the studies used the well-validated Nordic questionnaire while most of them relied on self-reporting of symptoms. Reporting of symptoms can have etiologies that range from non-organic to even serious pathologies. Multifactorial issues

influence self-perceived and self-reporting of symptoms among DP, such as their attitudes toward their profession,[45] work profile,[46] career choice,[47] career influences, levels of satisfaction with their work position, work stress,[48] salary, workplace preferences,[49] interest in community participation,[50] personal habits[51] etc. Other limitation noticed in the included studies was the interchangeable use of terms such as MSK disorders/ symptoms/pain. As healthcare professionals, pain is only one of the symptoms of MSK dysfunction. Other symptoms can be stiffness, discomfort, clicks, fatigue, etc which were often not reported in the earlier studies. Previous diagnosis of MSK disorders (specific and non-specific) by registered physicians was not observed in our reviewed studies as prevalence of MSK disorders.

Future studies can be done on evaluating the prevalence of MSK pain among DP in India, which surprisingly has not been reported before till date. Our Indian dentists and dental professionals are also prone to all the reviewed risk factors, and appropriate individualized and group ergonomic educational interventions might be indicated to ensure adequate prevention of MSK disorders. Outcome of such interventions could be evaluated using posture assessment instrument[52] validated for its use among dental operators.

CONCLUSION

There was an overall very high prevalence of MSK pain among DP. Among DP, dental hygienists and dental students were more affected with MSK symptoms. The risk factors highly predictive of developing MSK pain were biopsychosocial ,and further research on prevalence in Indian dental population is warranted before implementation of preventive educational programmes among dentists.

ACKNOWLEDGEMENTS

The authors wish to thank the authors of the reviewed studies for providing helpful

comments and valuable suggestions during the manuscript preparation process.

REFERENCES

1. Cruz GD, Ostroff JS, Kumar JV, Gajendra S. Preventing and detecting oral cancer- oral health care providers' readiness to provide health behavior counseling and oral cancer examinations. *J Am Dent Assoc* 2005; 136: 594-601.
2. Nermin Y. Musculoskeletal disorders (Msds) and dental practice: part-1. General information-terminology, aetiology, work-relatedness, magnitude of the problem and prevention. *Int Dent J* 2006; 56: 359-366.
3. Jacobsen N, Derand T, Hensten-Pettersen A. Profile of work-related health complaints among Swedish dental laboratory technicians. *Community Dent Oral Epidemiol* 1996; 24: 138-144.
4. Murtomaa H. Work-related complaints of dentists and dental assistants. *Int Arch Occup Environ Health* 1982; 50: 231-236.
5. Cimrin A, Komus N, Karaman C, Tertemiz KC. Pneumoconiosis and work-related health complaints in Turkish dental laboratory workers. *Tuberk Toraks* 2009; 57: 282-288.
6. Allsopp J, Basu MK, Browne RM, Burge PS, Matthews JB. Survey of the use of personal protective equipment and prevalence of work-related symptoms among dental staff. *Occup Environ Med* 1997; 54: 125-134.
7. Kanerva L, Alanko K, Jolanki R, Kanervo K, Susitaival P, Estlander T. The dental face mask- the most common cause of work-related face dermatitis in dental nurses. *Contact Dermatitis* 2001; 44 :261-262.
8. Szymanska J. Work-related noise hazards in the dental surgery. *Ann Agric Environ Med* 2000; 7: 67-70.
9. Szymanska J. Work-related vision hazards in the dental office. *Ann Agric Environ Med* 2000; 7:1-4.
10. Dajpratham P, Ploypetch T, Kiattavorncharoen S, Boonsiriseth K. Prevalence and associated factors of musculoskeletal pain among the dental personnel in a dental school. *J Med Assoc Thai* 2010; 93: 714-721.
11. Hayes MJ, Smith DR, Cockrell D. A systematic review of musculoskeletal disorders among dental professionals. *Int J Dent Hyg* 2009b; 7: 159-165.
12. Sartorio F, Vercelli S, Ferriero G, D'Angelo F, Migliario M, Franchignoni M. Work-related musculoskeletal diseases in dental professionals: 1- prevalence and risk factors. *G Ital Med Lav Ergon* 2005; 27: 165-169.
13. Lalumandier JA, McPhee SD, Parrott CB, Vendemia M. Musculoskeletal pain: prevalence, prevention and differences among dental office personnel. *Gen Dent* 2001; 49: 160-166.
14. Anton D, Rosecrance J, Merlino L, Cook T. Prevalence of musculoskeletal symptoms and carpal tunnel syndrome among dental hygienists. *Am J Ind Med* 2002; 42: 248-257.
15. Osborn JB, Newell KJ, Rudney JD, Stoltenberg JL. Musculoskeletal pain among Minnesota dental hygienists. *J Dent Hyg* 1990; 64: 132-138.
16. de Carvalho MV, Soriano EP, de Franca Caldas A Jr, Campello RI, de Miranda HF, Cavalcanti FI. Work-related musculoskeletal disorders among Brazilian dental students. *J Dent Educ* 2009; 73: 624-630.
17. Thornton LJ, Barr AE, Stuart-Buttle C, Gaughan JP, Wilson ER, Jackson AD et al. Perceived musculoskeletal symptoms among dental students in the clinic environment. *Ergonomics* 2008; 51: 573-586.
18. Abou-Atme YS, Melis M, Zawawi KH, Cottogno L. Five-year follow-up of temporomandibular disorders and other musculoskeletal symptoms in dental students. *Minerva Stomatol* 2007; 56: 603-609.
19. Tezel A, Kavrut F, Tezel A, Kara C, Demir T, Kavrut R. Musculoskeletal disorders in left- and right- handed Turkish dental students. *Int J Neurosci* 2005; 115: 255-266.
20. Melis M, Abou-Atme YS, Cottogno L, Pittau R. Upper body musculoskeletal symptoms in Sardinian dental students. *J Can Dent Assoc* 2004; 70: 306-310.
21. Rice VJ, NindlB, Pentikis JS. Dental workers, musculoskeletal cumulative trauma and carpal tunnel syndrome, who is at risk? A pilot study. *Int J Occup Saf Ergon* 1996; 2: 218-233.
22. Hayes MJ, Smith DR, Cockrell D. Prevalence and correlates of musculoskeletal disorders among Australian dental hygiene students. *Int J Dent Hyg* 2009a; 7: 176-181.
23. Samotoi A, Moffat SM, Thomson WM. Musculoskeletal symptoms in New Zealand dental therapists: prevalence and associated disability. *N Z Dent J* 2008; 104: 49-53.
24. Finsen L, Christensen H, Bakke M. Musculoskeletal disorders among dentists and variation in dental work. *Appl Ergon* 1998; 29: 119-125.
25. Augustson TE, Morken T. Musculoskeletal problems among dental health personnel- a survey of the public dental health services in Hordaland. *Tidsskr Nor Laegeforen* 1996; 116: 2776-2780.

26. Oberg T, Oberg U. Musculoskeletal complaints in dental hygiene: a survey study from a Swedish country. *J Dent Hyg* 1993; 67: 257-261.
27. Morse T, Bruneau H, Dussetschleger J. Musculoskeletal disorders of the neck and shoulder in the dental professions. *Work* 2010; 35: 429-429.
28. Morse T, Bruneau H, Michalak-Turcotte C, Sanders M, Warren N, Dussetschleger J et al. Musculoskeletal disorders of the neck and shoulder in dental hygienists and dental hygiene students. *J Dent Hyg* 2007; 81: 10.
29. Akesson I, Johnsson B, Rylander L, Moritz U, Skerfving S. Musculoskeletal disorders among female dental personnel- clinical examination and a 5-year follow-up study of symptoms. *Int Arch Occup Environ Health* 1999; 72: 395-403.
30. Kihara T. Dental care works and work-related complaints of dentists. *Kurume Med J* 1995; 42: 252-257.
31. Valachi B. Musculoskeletal health of the woman dentist: distinctive interventions for a growing population. *J Calif Dent Assoc* 2008; 36: 127-132.
32. Yamalik N. Musculoskeletal disorders (MSDs) and dental practice part-2- risk factors for dentistry, magnitude of the problem, prevention and dental ergonomics. *Int Dent J* 2007; 57: 45-54.
33. Ylipaa V, Szuster F, Spencer J, Preber H, Benko SS, Arnetz BB. Health, mental well-being and musculoskeletal disorders: a comparison between Swedish and Australian dental hygienist. *J Dent Hyg* 2002; 76: 47-58.
34. Akesson I, Schutz A, Horstmann V, Skerfving S, Moritz U. Musculoskeletal symptoms among dental personnel: lack of association with mercury, selenium status, overweight and smoking. *Swed Dent J* 2000; 24: 23-38.
35. Moen BE, Bjorvatn K. Musculoskeletal symptoms among dentists in a dental school. *Occup Med* 1996; 46: 65-68.
36. Locker D, Burman D, Otchere D. Work-related stress and its predictors among Canadian dental assistants. *Community Dent Oral Epidemiol* 1989; 17: 263-266.
37. Warren N. Causes of musculoskeletal disorders in dental hygienists and dental hygiene students: a study of combined biomechanical and psychosocial risk factors. *Work* 2010; 35: 441-454.
38. Ylipaa V, Arnetz BB, Preber H. Predictors of good general health, well-being and musculoskeletal disorders in Swedish dental hygienists. *Acta Odontol Scand* 1999; 57: 277-282.
39. More TF, Michalak-Turcotte C, Atwood-Sanders M, Warren N, Peterson DR, Bruneau H et al. A pilot study of hand and arm musculoskeletal disorders in dental hygiene students. *J Dent Hyg* 2003; 77: 173-179.
40. Ylipaa V, Anetz BB, Benko SS, Ryden H. Physical and psychosocial work environments among Swedish dental hygienists: risk indicators for musculoskeletal complaints. *Swed Dent J* 1997; 21: 111-120.
41. Liss GM, Jesin E, Kusiak RA, White P. Musculoskeletal problems among Ontario dental hygienists. *Am J Ind Med* 1995; 28: 521-540.
42. Michalak-Turcotte C. Controlling dental hygiene work-related musculoskeletal disorders: the ergonomic process. *J Dent Hyg* 2000; 74: 41-48.
43. Sanders MA, Turcotte CM. Strategies to reduce work-related musculoskeletal disorders in dental hygienists: two case studies. *J Hand Ther* 2002; 15: 363-374.
44. Karjalainen KA, Malmivaara A, van Tulder MW, Roine R, Jauhiainen M, Hurri H, Koes BW. Multidisciplinary biopsychosocial rehabilitation for neck and shoulder pain among working age adults. *Cochrane Database Syst Rev* 2003; 2: CD002194.
45. Saito A, Tomita C, Sato Y, Cathcart G. Perceptions of Japanese and Canadian dental hygiene students towards their profession. *Int J Dent Hygiene* 2009; 7: 188-195.
46. Tseveenjev B, Virtanen JI, Wang NJ, Widstrom E. Working profiles of dental hygienists in public and private practice in Finland and Norway. *Int J Dent Hygiene* 2009; 7: 17-22.
47. DeAngelis S, Dean K, Pace C. Career choice and perceptions of dental hygiene students and applicants. *J Dent Hyg* 2003; 77: 97-105.
48. Gorter RC. Work stress and burnout among dental hygienists. *Int J Dent Hygiene* 2005; 3: 88-92.
49. Petre'n V, Levin G, Chohan T, Preber H, Candell A, Bergstro'm J. Swedish dental hygienists' preferences for workplace improvement and continuing professional development. *Int J Dent Hygiene* 2005; 3: 117-125.
50. Bhayat A, Yengopal V, Rudolph MJ, Naidoo U, Vayej A. Attitudes of South African oral hygienists towards compulsory community service. *Int J Dent Hygiene* 2008; 6: 8-12.
51. Bergstrom J, Petre'n V, Bark J, Preber H. Smoking habits among Swedish dental hygienists: a 15-year perspective (1987-2002). *Int J Dent Hygiene* 2009; 7: 49-54.
52. Branson BG, Williams KB, Kimberly KB, McInay SL, Dickey D. Validity and reliability of a dental operator posture assessment instrument. *J Dent Hyg* 2002; 76: 255-262.