

SHORT COMMUNICATION

Use of Ocular Biomarkers for Predicting Systemic Bone Health in Orthopaedic Patients

Jeff Walter Rajadurai OR¹, Radhika Dinesh², S. Jeeva³

HOW TO CITE THIS ARTICLE:

Jeff Walter Rajadurai OR, Radhika Dinesh, S. Jeeva. Use of Ocular Biomarkers for Predicting Systemic Bone Health in Orthopaedic Patients. *Jr. Orth. Edu.* 2025; 11(2): 65–68.

ABSTRACT

Using signs from the eye to understand what's going on in the bones might sound unusual at first but it's a line of thought that's slowly gaining momentum. After all, a number of systemic diseases show up in the eye, and doctors have long relied on this connection. So, it's not a big leap to consider whether eye health could also reflect bone health, especially in conditions like osteoporosis. In this work, we take a closer look at whether certain patterns things like how blood vessels look at the back of the eye, how stiff the cornea is, or changes around the optic disc might offer early clues about bone weakness. It's not just guesswork; some of these features seem to share the same underlying causes as bone deterioration: inflammation, tiny vessel problems, and oxidative stress, to name a few. With tools like OCT and new ways to map the retinal vessels, clinicians might be able to pick up these signs well before bone loss becomes obvious. For orthopaedic doctors, that could mean acting earlier, perhaps even adjusting care plans based on what's seen in the eye. And maybe just maybe this could shift the focus a bit. From reacting to fractures to trying to prevent them. That's the bigger picture. Of course, a lot more research is needed. But the potential is there, and it's worth paying attention to.

KEYWORDS

• Biomarkers • Osteoporosis • Optical Coherence Tomography • Retinal Microvasculature • Corneal Biomechanics

AUTHOR'S AFFILIATION:

¹ Associate Professor, Department of Orthopedics, Madha Medical College & Research Institute, Chennai, Tamil Nadu, India.

² Assistant Professor, Ophthalmology, ESIC Medical College, KK Nagar, Chennai, Tamil Nadu, India.

³ Assistant Professor, Meenakshi Medical College and Research Institute, Meenakshi Nagar, Tamil Nadu, India.

CORRESPONDING AUTHOR:

Jeff Walter Rajadurai OR, Associate Professor, Department of Orthopedics, Madha Medical College & Research Institute, Chennai, Tamil Nadu, India.

E-mail: jeffy.walter@gmail.com

➤ Received: 21-03-2025 ➤ Accepted: 03-06-2025



Creative commons non-commercial CC BY-NC: This article is distributed under the terms of the creative commons attribution non-commercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the Red Flower Publication and Open Access pages (<https://rfppl.co.in>)

INTRODUCTION

The human body is essentially an interconnected system with the cooperation of many organ systems to preserve homeostasis. In this regard, there is an increasing interest in the connection between systemic bone problems and eye health. In the aging populations, osteoporosis and other metabolic bone problems are becoming very common, sometimes remaining undiagnosed until fractures or severe effects occur. This study, therefore, suggests that easily accessible, non-invasive ocular biomarkers may be used as a surrogate for systemic bone health and may offer doctors with essential diagnostic information.

This multimodal approach makes use of advances in ocular imaging and genetic diagnostics to highlight the connections between bone and eye health. The vascular network of the retina, corneal biomechanics, and optic nerve anatomy provide a window into systemic processes. This study suggests the incorporation of ocular biomarkers into orthopaedic diagnostic frameworks and explaining how they predict systemic bone health to improve patient outcomes.

Ocular Biomarkers and Systemic Bone Health: A Cross-Sectional Analysis

1. Retinal Microvasculature and Bone Density

The microvascular network of the retina is a useful indicator of overall health because it is highly responsive to systemic changes. Research indicates that individuals with low bone mineral density (BMD) often have reduced retinal vascular density and increased vascular tortuosity. Unlike isolated alterations, these are indicative of metabolic bone diseases that lead to systemic microvascular dysfunction. These microvascular changes can be seen in great detail because of non-invasive retinal imaging, particularly optical coherence tomography angiography (OCTA). The high-resolution imaging capabilities of OCTA allow clinicians to quantitatively correlate microvascular health with systemic bone health markers. By recognising these trends, orthopaedic specialists can determine an individual's risk of fractures and other issues associated with osteoporosis.^{1,2}

2. Corneal Biomechanics as a Marker of Systemic Health

The biomechanical properties of the cornea are another critical indicator of systemic health.

Measurement of the corneal resistance factor and corneal hysteresis, which quantify the cornea's viscoelastic properties, is altered by systemic diseases such as osteoporosis. Osteoporotic patients often have lower levels of corneal stiffness, which may indicate that loss of collagen is a common mechanism of pathophysiology. Devices such as the ORA are increasingly used for the evaluation of these attributes. These methods provide orthopaedic doctors with a useful but indirect view of how important systemic collagen integrity is to bone and ocular health. The non-invasive assessment of ocular biomechanics opens up the diagnostic toolbox for orthopaedic practitioners.³

3. Optic Disc Morphology and Bone Health

Further information concerning the health of systemic bones can be gathered from the changes in morphology of the optic disc. Increased optic cupping and thinning of the RNFL are associated with systemic diseases such as osteoporosis. These changes are thought to be due to vascular abnormalities and redistribution of mechanical stress, which deteriorate the fragility of bones. Advanced imaging methods, Heidelberg Retinal Tomography (HRT), in particular, would detect minor but significant optic disc configuration changes. Systemic assessments of bone density might possibly be compared against these discoveries to improve diagnostic processes. Patients at risk of severe osteoporosis would thus receive targeted and timely interventions.^{4,5}

Shared Pathophysiological Mechanisms

1. Oxidative Stress and Inflammation

Oxidative stress and chronic inflammation are essential factors in the preservation of bone and eye health. Oxidative stress is responsible for microvascular damage and degenerative retinal disorders. Similarly, oxidative stress increases osteoclastic activity and decreases osteoblastic function in bones leading to decreased bone density. Chronic inflammation worsens this impact because of the release of pro-inflammatory cytokines that cause more damage to the skeletal and circulatory tissues. With these pathways being targeted, treatments such as antioxidants and anti-inflammatory drugs can potentially improve bone as well as eye health simultaneously.⁶

2. Hormonal Interactions

Hormonal regulation is another important link between systemic and ocular bone health. Oestrogen, a hormone crucial to maintaining bone density, also maintains the tissues of the eyes by regulating collagen and vascular health. The simultaneous decline in bone density and ocular integrity in postmenopausal women underlines the hormonal connection. A more holistic approach to diagnosis and treatment of systemic bone issues may be provided by hormonal evaluations in conjunction with ocular biomarker research.⁷

3. Vascular Dysregulation

The health of the circulatory system is critical for both skeletal and ocular health. Retinal vascular abnormalities that are indicative of more serious systemic vascular issues include microaneurysms and capillary dropout. These vascular abnormalities may disrupt the flow of nutrients, affecting the development of the retina and bone remodelling. Therefore, retinal imaging is a useful indicator of systemic vascular health and provides information about underlying bone health concerns.⁸

Technological Advances in Ocular Biomarker Detection

1. Optical Coherence Tomography

OCT has become a new gold standard in ocular diagnostics, providing high-resolution images of retinal and optic nerve structures. Its sensitivity in detecting subtle changes in thickness of the retinal layer and microvascular integrity allows it to be used primarily in identifying early signs of systemic bone health deterioration. In orthopaedic patients, OCT can supplement traditional diagnostic tools by providing another layer of understanding of systemic health.⁹

2. Corneal Imaging

ORA and CORVIS ST devices have taken up a new face to assessment in corneal biomechanics. It enables them to provide exact corneal viscoelastic and resistance measurements which, indeed is related to systemic integrity of collagen. Introducing corneal imaging in orthopaedic assessment may provide valuable insights into a patient's systemic health status.

3. Artificial Intelligence in Diagnosis

With AI in ocular diagnostics, it's easier to see hidden patterns and correlations that the

human mind cannot easily see. Algorithms using AI can go through immense amounts of data from OCT imaging and retinal vascular analysis, predicting systemic bone health. All this would help in raising diagnostic precision but also individualize treatment options.¹⁰

Clinical Implications and Future Directions

The management of systemic bone defects may be revolutionized through orthopaedic treatment combined with ocular biomarkers. With high technological imaging, the ordinary use of eye tests can serve as an early detection of osteoporosis and other similar conditions. Through the early detection before the crippling symptoms have actually occurred, this approach induces a shift in the response to treatment from being reactionary to being preventative.

Long-term studies to establish the predictive value of ocular biomarkers should be the focus of future research. In addition, the therapeutic outcomes of focused treatments targeting the condition of the ocular and systemic bones may lead to the establishment of new integrated care approaches. A multidisciplinary approach such as this ensures that patients receive complete care through encouraging cooperation between orthopaedic physicians and ophthalmologists.

CONCLUSION

Ocular biomarkers have revolutionised assessment of systemic bone health through their discovery. This method represents a low-cost, non-invasive means to provide early diagnosis and intervention while bridging the disciplines of orthopaedics and ophthalmology. As long as technology keeps redefining these tools, it promises to improve outcomes in patients and encourages cross-functional interactions.

Source(s) of Support and Funding: Nil

Conflict of Interest: Nil

Ethical Approval and/or Institutional Review Board (IRB): Not applicable

REFERENCES

1. Spaide R.F., Klancnik J.M., Cooney M.J. Retinal vascular imaging using optical coherence tomography angiography. *Retina*. 2015; 35(11): 2181-6.

2. Cheung C.Y., Ikram M.K., Sabanayagam C., Wong T.Y. Retinal microvasculature as a model to study the manifestations of hypertension. *Hypertension*. 2012; 60(5): 1094-103.
3. Wells A.P., Garway-Heath D.F., Poostchi A., Wong T., Gazzard G., Foster P.J. Corneal hysteresis as a risk factor for glaucoma progression: a prospective study. *Ophthalmology*. 2008; 115(9): 1501-5.
4. Frost S.A., Nguyen N.D., Center J.R., Eisman J.A., Nguyen T.V. Risk factors for hip fracture in men aged 50 years and older: a prospective study. *Bone*. 2013; 52(1): 419-26.
5. Bourne R.R., Stevens G.A., White R.A., *et al.* Causes of vision loss worldwide, 1990-2010: a systematic analysis. *Lancet Glob Health*. 2013; 1(6): e339-49.
6. Phipps J.A., Yaganti S., Schlotzer-Schrehardt U, *et al.* Proteomic markers of exfoliation syndrome support a chronic stress response in the trabecular meshwork. *Exp Eye Res*. 2012; 101: 27-34.
7. Raisz LG. Pathogenesis of osteoporosis: concepts, conflicts, and prospects. *J Clin Invest*. 2005; 115(12): 3318-25.
8. Keenan T.D., Goldacre R., Goldacre M.J. Associations between primary open angle glaucoma, Alzheimer's disease and vascular comorbidities: record linkage study. *Br J Ophthalmol*. 2015; 99(5): 665-8.
9. Schuman J.S., Hee M.R., Puliafito C.A., *et al.* Quantification of nerve fiber layer thickness in normal and glaucomatous eyes using optical coherence tomography. *Arch Ophthalmol*. 1995; 113(5): 586-96.
10. Gulshan V., Peng L., Coram M., *et al.* Development and validation of a deep learning algorithm for detection of diabetic retinopathy in retinal fundus photographs. *JAMA*. 2016; 316(22): 2402-10.